



International Journal for Innovative Engineering and Management Research

A Peer Reviewed Open Access International Journal

www.ijiemr.org

COPY RIGHT

2020 IJIEMR. Personal use of this material is permitted. Permission from IJIEMR must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collective works, for resale or redistribution to servers or lists, or reuse of any copyrighted component of this work in other works. No Reprint should be done to this paper, all copy right is authenticated to Paper Authors

IJIEMR Transactions, online available on 24th Mar 2020. Link

[:http://www.ijiemr.org/downloads.php?vol=Volume-09&issue=ISSUE-03](http://www.ijiemr.org/downloads.php?vol=Volume-09&issue=ISSUE-03)

Title: **SPECTROPHOTOMETRIC DETERMINATION OF MEROPENEM IN BULK AND INJECTION FORMULATIONS BY BRUCINE**

Volume 09, Issue 03, Pages: 89-95.

Paper Authors

DR ARUNA KUMARI NAKKELLA, PROF. K. RAGHU BABU, DR P SRIRAMA MURTHY



USE THIS BARCODE TO ACCESS YOUR ONLINE PAPER

To Secure Your Paper As Per **UGC Guidelines** We Are Providing A Electronic Bar Code

SPECTROPHOTOMETRIC DETERMINATION OF MEROPENEM IN BULK AND INJECTION FORMULATIONS BY BRUCINE

¹DR ARUNA KUMARI NAKKELLA, ²PROF. K. RAGHU BABU, ³DR P SRIRAMA MURTHY

¹Asst Principal, College of Engineering, Dr B R Ambedkar University Etcherla, Srikakulam

²Registrar, Dr B R Ambedkar University, Etcherla, Srikakulam

³Principal (Retired), SKVT Degree and PG College Rajahmundry

drsiv@gmail.com

ABSTRACT

A simple and cost effective spectrophotometric method was described for the determination of Meropenem in pure form and in pharmaceutical formulations. The method is based on the formation of colored chromogen when the drug reacts with brucine and sodium periodate in acidic medium. This method was applied for the determination of drug contents in pharmaceutical formulations and enabled the determination of the selected drug in microgram quantities (0.5 to 3.0 mL). No interferences were observed from excipients and the validity of the method was tested against reference method. The colored species has an absorption maximum at 520 nm for Meropenem and obeys Beer's law in the concentration range 0.02-0.12 mg/mL of Meropenem. The apparent molar absorptivity was 61×10^5 and Sandell's sensitivity was 7×10^{-4} . The slope is 0.0230 ± 0.0008 , the intercept of the equation of the regression line is 0.0230 ± 0.0008 . The optimum experimental parameters for the reaction have been studied and the validity of the described procedure was assessed. Statistical analysis of the results has been carried out revealing high accuracy and good precision. The proposed method was successfully applied for the determination of Meropenem in pharmaceutical formulations.

Keywords- Meropenem, Brucine, sodium periodate, Spectrophotometry.

INTRODUCTION

Due to counterfeiting, the drug quality has become a source of major concern worldwide, particularly in many developing countries. The most commonly counterfeited drugs are anti-infectives or antibiotics. Use of poor quality antibiotics bears serious health implications such as treatment failure, adverse reactions, drug resistance, increased morbidity, and mortality¹. Among antibiotics, penems are much recently

introduced, widely prescribed and costlier. Therefore, incentive to produce their counterfeits because of profit margin increases considerably.

Meropenem² is a broad spectrum beta-lactam antibiotic belonging to the carbapenem class.

1.1 Drug Profile

Name : Meropenem (MRP)
 Chemical Name : (3-[5-(dimethylcarbamoyl)pyrrolidin-2-yl]sulfanyl-6-(1-hydroxyethyl)-4-methyl-7-oxo-1-azabicyclo[3.2.0]hept-2-ene-2-carboxylic acid




Fig: 2.2.1

Molecular formula : $C_{12}H_{17}N_3O_4S$
 Molecular weight : 383.46 g/mol
 Empirical formula : $C_{12}H_{17}N_3O_4S \cdot H_2O$
 Color : Off-white
 pK_a : 3.47
 Solubility : Sparingly soluble in water and slightly soluble in methanol
 Pharmacodynamic / Chemotherapeutic category : Antibacterial Agent

Meropenem acts by interfering with their ability to form cell walls, and therefore the bacteria break up and die. It is a broad spectrum antibiotic with activity against many aerobic and anaerobic gram-positive and gram-negative organisms. In contrast to other beta-lactams, it is highly resistant to degradation by beta-lactamases or cephalosporinases.

Literature survey reveals that the drugs were determined by using HPLC and some spectrophotometric methods for Meropenem³⁻⁸. According to literature survey there is no method reported for Meropenem with Brucine reagent by visible spectrophotometry. Hence an attempt was made to develop sMRPle and sensitive spectrophotometric method for the estimation of the above drug in pure and in

pharmaceutical formulations. The method uses the well known oxidative coupling reaction between the reagent and Meropenem resulting in the formation of a coloured chromogen that could be measured at 520 nm for Meropenem.

2. EXPERIMENTAL

2.1 Apparatus

All spectral characteristics and absorbance measurements were made on Perkin Elmer, LAMBDA 25 double beam UV-Visible spectrophotometer with 10 mm matched quartz cells. All chemicals used were of analytical reagent grade and double distilled water was used throughout. Brucine supplied by Loba chemicals Ltd., India, was used by diluting 200.0mg to 100 mL with distilled water. $NaIO_4$ supplied by BDH chemicals Ltd., India, was used by dissolving 200.0 mg of Sodium meta periodate to 100 mL with distilled water. H_2SO_4 (18 M) supplied by Qualigens was used as it is. 10 mg/mL stock reference solution was freshly prepared from pure sample of Meropenem by dissolving 100 mg in 100 mL of double distilled water.

2.2 General procedure

Method

Into a series of 10 mL volumetric flask, different aliquots of working standard solution (0.5 to 3.0 mL) of MRP were transferred to provide final concentration range of 0.02 – 0.12 mg/mL. To each flask, 3 mL of BCN(0.2%) , 2 mL of $NaIO_4$ (0.2%) and 2.5 mL of H_2SO_4 were successively added and kept aside for 5 min. The solutions were made up to volume with


distilled water. The absorbance of each solution was measured at 523 nm against the reagent blank. The calibration graph was then prepared by plotting the absorbance versus the concentration of the drug. The concentration of the unknown was read from the calibration graph or computed from the regression equation.

2.3 Procedure for Injections

An amount of powder equivalent to 100 mg of Meropenem was weighed into a 100 mL volumetric flask, 50 mL of distilled water was added and shaken thoroughly for about 10 min, then the volume was made up to the mark with the distilled water, mixed well and filtered. Further dilutions were made and the assay of injections was completed according to general procedure.

3. RESULTS AND DISCUSSION

The dimethoxy benzene nucleus of Brucine is attacked by IO_4^- with the formation of o-quinone (bruciquinone) which in turn undergoes nucleophilic attack on the most electron rich position of the coupler i.e., proton bearing amino group (secondary amine in MRP), to give 1-mono substituted bruciquinone derivative. In the present study, MRP was successfully determined by this method. The reactions of the MRP with brucine in the presence of IO_4^- are described in the Scheme - 1.



Scheme - 1

4. Optimization of conditions on absorption spectrum of the reaction product

The condition under which the reaction of Meropenem with Brucine fulfills the essential requirements was investigated. All conditions studied were optimized at room temperature ($32 \pm 2^\circ\text{C}$).

4.1 Selection of reaction medium

To find a suitable medium for the reaction, different acidic mediums can be used. The best results were obtained when H_2SO_4 was used. In order to determine the optimum concentration of H_2SO_4 , different volumes of H_2SO_4 solution (0.5 – 3.0 mL) were used to a constant concentration of Meropenem (1mg/mL) and the results were observed. From the absorption spectrum it was evident that 2.0 mL of H_2SO_4 solution was found optimum. Larger volumes had no significant effect on the absorbance of the colored species.

4.2 Effect of order of addition of reactants

Few trials were performed to ascertain the influence of order of addition of reactants on the color development and the results are presented in Table 1. The order of addition

of serial number (i) is recommended for

Table 1. Effect of order of addition of reactants on color development.

S.No.	Drug	Order of Addition	Absorbance	Recommended order of Addition
1.	Meropenem ^a	i D+ BCN+NaIO ₄	0.189	i
		ii D+ NaIO ₄ +BCN	0.04	
		iii BCN+ NaIO ₄ +D	0.145	

^aFor 40 µg/mL of Drug samples

Meropenem.

4.3 Effect of Brucine concentration

Several experiments were carried out to study the influence of BCN concentration on the color development by keeping the concentration of drug and Sodium meta periodate to constant and changing reagent concentration. It was apparent that 2.5 mL of BCN gave maximum color for Meropenem

4.4 Effect of Sodium meta periodate concentration

Several experiments were carried out to study the influence of Sodium meta periodate concentration on the color development by keeping the concentration of drug, BCN to constant and changing Sodium meta periodate concentration. It was apparent that 2.5 mL of reagent gave maximum color for Meropenem.

5. REACTION TIME AND STABILITY OF THE COLORED SPECIES

The color reaction was not instantaneous. Maximum color was developed within 5 minutes of mixing the reactants and was stable for 60 minutes thereafter.

Tab: 2.6.2 RESULTS OF METHOD OPTIMISATION FOR MRP – BCN

Parameter	Range of study	Optimised condition in procedure	Remarks
λ_{max} (nm)	400-600	523	
Effect of volume of BCN required for oxidative coupling (mL)	0.5-4.0	3.0	Volume above 3.0 mL gave high optical densities in blanks (>3.0), which resulted in deviations from Beers law.
Effect of volume of H ₂ SO ₄ (mL)	0.5-3.0	2.5	To speed up the reaction in color development, 2.5 mL of H ₂ SO ₄ (18M) was found necessary for maximum color development.
Effect of volume of NaIO ₄ (mL)	0.5-3.0	2.0	Addition of 2.0 mL of NaIO ₄ was found optimum for oxidizing the drug.
Effect of reaction Time (min)	15-30	15	The minimum time required for complete oxidation was found to be 15 min.
Effect of temperature (°C) for oxidative coupling	20-40	32 ± 2 Lab. Temp	At low temperatures (<30°C) the reaction time was found to be more and at high temperatures (>34°C) no added advantage was found.
Standing time (min)	1-3	2	A minimum amount of time, i.e., 2 min was necessary for undergoing oxidative coupling and beyond 3 min results in low sensitivity.
Stability period after final dilution (min)	5-40 min	40 min	The absorbance of the colored product decreases slowly with time after 40 min.

6. ABSORPTION SPECTRUM AND CALIBRATION GRAPH

Absorption spectrum of the colored complex was scanned at 450-850 nm against a reagent blank. The reaction product showed absorption

maximum at 520 nm for Meropenem. Calibration graph was obtained according to the above general procedure. The linearity replicates for six different concentration of Meropenem was checked by a linear least - squares treatment. All the spectral characteristics and the measured or calculated factors and parameters were summarized in Table 2.

Fig 1. Calibration graph of Meropenem





Fig 2 Absorption spectra of Meropenem



6.1 Sensitivity, accuracy and precision

Sandell's sensitivity, molar absorptivity, precision and accuracy were found by performing eight replicate determinations containing 3/4th of the amount of upper Beer's law limits. The measured standard deviation (S.D), relative standard deviation (RSD), and confidence limits (Table 2) were considered satisfactory.

6.2 Interference

These substances are seldom present in the reagents and used in the pharmaceutical formulations. Hence, the method is devoid of error due to above substances.

Table 2. Optical and regression characteristics of the proposed method for Meropenem.

λ_{max} nm	520	
Bear's law limits, $\mu\text{g/mL}$	0.02-0.12	
Molar absorptivity, L/mol.cm	61×10^4	
Sandell's sensitivity $\mu\text{g/cm}^2/0.001$ absorbance unit	7×10^4	
Regression equation ($Y = a + bc$)		
Slope(b)	0.0230 ± 0.0008	
Standard deviation of slope (Sb)	0.0021	
Intercept	0.0029 ± 0.0014	
r^2	0.9939	
Limit of Detection	0.00667	
Limit of Quantification	0.0998	
Standard deviation of intercept (Sa)	0.0023	
Standard error of estimation (Se)	0.0121	
Correlation coefficient @	0.9998	
Relative standard deviation (%)*	0.0480	
% Range of error (Confidence limits)*		
Precision		
0.05 level	0.2231	
0.01 level	0.3196	
Accuracy		
Bulk sample	Amount found (μg)	% error
50	49.54	0.92
75	74.95	0.06
100	99.54	0.46

7. APPLICATION TO FORMULATION

Table 3. Results of analysis of injection formulations containing Meropenem

Injection	Meropenem
Company Name	Troika Pharma
Formulation	Inj
Labeled amount, mg	1000
% Recovery	99.8

8. CONCLUSION

The proposed method was found to be sMRPle, rapid and inexpensive, hence can be used for routine analysis of Meropenem in bulk and in injection formulations.

9. ACKNOWLEDGEMENT

We wish to thank Aurobindo labs, Hyd. for providing gifted samples of Penems; Research lab, Dept., of Engineering chemistry, AUCE(A), Visakhapatnam,

India, Dept., of Analysis, GIET School of Pharmacy, Rajahmundry, India.

10. REFERENCES

1. United States Pharmacopeia Drug Quality and Information Program. 2004. A review of drug quality in Asia with focus on anti-infectives, United States Pharmacopoeia, Drug Quality and Information Program 1-46.



2. Sean C. Sweetman, Martindale Extra Pharmacopoeia 36(1), 286, Pharmaceutical Press, 2009.
3. Forsyth R J and Ip DP, J Pharm Biomed Anal, "Determination of Meropenem and Cilastatin sodium in Primaxin by first order derivative ultraviolet spectrophotometry", 12(10), 1243-8, 1994.
4. Gravallesse D A, Musson D G, Pauliukonis L T, Bayne W F., "Determination of Meropenem (N-formimidoylthienamycin) in human plasma and urine by high-performance liquid chromatography, comparison with microbiological methodology and stability", J Chromatography, 14(1), 71-84, 1984.
5. Myers C M and J L Blumer J L, "Determination of Meropenem and Cilastatin in serum by high-pressure liquid chromatography", Antimicrob Agents Chemother, 26(1), 78-81, 1984.
6. Garcia- Capdevila L, López-Calull C, Arroyo C, Moral M A, Mangues M A and Bonal J, "Determination of Meropenem in plasma by high-performance liquid chromatography for pharmacokinetic studies in patients", J Chromatogr B Biomed Sci Appl 25(1), 127-132, 1997.
7. Irene A, Miguel A B, Manuel C, and Juan C J, "Liquid chromatographic method for the simultaneous determination of Meropenem and sulbactam in Mouse Plasma". J. chromatography Sci. 44, 548-551, 2006.
8. Chaudhary A K, Ankushrao W S, Yadav S, Chandrashekhar T G and Vandana S, "Validated Reverse Phase HPLC method for the determination of DEHP content in reconstituting diluents and in reconstituted solutions of Meropenem and Cilastatin for Injection", E-J. Chem., 7(2), 501-513, 2010.