International Journal for Innovative Engineering and Management Research
PEER REVIEWED OPEN ACCESS INTERNATIONAL JOURNAL
www.ijiemr.org

COPY RIGHT



ELSEVIER SSRN

2024 IJIEMR. Personal use of this material is permitted. Permission from IJIEMR must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating newcollective works, for resale or redistribution to servers or lists, or reuse of any copyrighted component of this work in other works. No Reprint should be done to this paper; all copy right is authenticated to Paper Authors

IJIEMR Transactions, online available on 4th April 2024. Link

https://ijiemr.org/downloads.php?vol=Volume-13&issue=issue04

DOI:10.48047/IJIEMR/V13/ISSUE04/78

Title: " FORMULATION INNOVATIONS: BOOSTING DISSOLUTION OF HERBAL BIOACTIVES VIA NATURAL POLYMERS AND PHYTOCONSTITUENT INTERACTIONS"

Volume 13, ISSUE 04, Pages: 680-685

Paper Authors
Mallikarjun N. Guddin, Dr. Shiv Kumar Gupta

USE THIS BARCODE TO ACCESS YOUR ONLINE PAPER

To Secure Your Paper as Per UGC Guidelines We Are Providing A ElectronicBar code

International Journal for Innovative Engineering and Management Research
PEER REVIEWED OPEN ACCESS INTERNATIONAL JOURNAL
www.ijiemr.org

FORMULATION INNOVATIONS: BOOSTING DISSOLUTION OF HERBAL BIOACTIVES VIA NATURAL POLYMERS AND PHYTOCONSTITUENT INTERACTIONS

[1]Mallikarjun N. Guddin, [2]Dr. Shiv Kumar Gupta (Associate Professor)
[1]Research Scholar, Glocal School of Pharmacy,The Glocal University, Mirzapur Pole, Saharanpur, U.P.
[2]Supervisor, Glocal School of Pharmacy,The Glocal University, Mirzapur Pole, Saharanpur, U.P.

ABSTRACT

Herbal bioactives offer immense therapeutic potential; however, their clinical application is frequently hampered by poor aqueous solubility and erratic bioavailability. This study explores formulation innovations that leverage natural polymers to enhance the dissolution of herbal bioactives through strategic interactions with phytoconstituents. By employing various approaches—including solid dispersions, nanoparticulate systems, and inclusion complexation—we demonstrate that natural polymers not only improve the wettability and dissolution rate of these compounds but also protect them from environmental degradation. In vitro dissolution studies reveal a marked improvement in the solubility profiles of selected herbal actives, suggesting that biopolymer-phytoconstituent interactions could serve as a promising platform for developing effective and sustainable herbal formulations.

Keywords: Natural polymers, herbal bioactives, dissolution enhancement, phytoconstituents, drug delivery

I. INTRODUCTION

Herbal bioactives have long been celebrated for their multifaceted pharmacological properties, ranging from anti-inflammatory and antioxidant effects to anticancer and antimicrobial activities. Despite these therapeutic potentials, the clinical utility of many plant-derived compounds is significantly limited by their inherent poor aqueous solubility and low oral bioavailability. This solubility challenge often results in reduced therapeutic efficacy, necessitating high dosages that can inadvertently lead to adverse side effects.

Natural polymers, such as chitosan, alginate, cellulose derivatives, and cyclodextrins, have emerged as effective carriers to improve the solubility and delivery of hydrophobic herbal compounds. Their biocompatibility, biodegradability, and functional versatility allow for the creation of innovative formulation strategies that optimize drug dissolution rates and sustain release profiles. The interaction between these polymers and phytoconstituents can lead to the formation of amorphous solid dispersions, nanoparticles, and inclusion complexes, all of which contribute to enhanced solubility and stability of the active ingredients.

This paper investigates the formulation innovations that utilize natural polymers to boost the dissolution of herbal bioactives. By examining the mechanisms of polymer-phytoconstituent

interactions, we aim to provide a comprehensive overview of the current strategies and their potential to revolutionize herbal drug delivery systems.

II. MATERIALS AND METHODS

In this study, we selected three model herbal bioactives—curcumin, quercetin, and resveratrol—known for their poor aqueous solubility. Natural polymers including chitosan, hydroxypropyl methylcellulose (HPMC), and β-cyclodextrin were employed as solubilizing agents.

Preparation of Formulations

Solid Dispersions:

Herbal bioactives were dispersed in a hydrophilic polymer matrix (HPMC) using the solvent evaporation method. Briefly, the bioactive and polymer were dissolved in ethanol, and the solvent was subsequently evaporated under reduced pressure to yield a uniform amorphous dispersion.

Nanoparticulate Systems:

Chitosan nanoparticles were prepared using the ionic gelation method. The bioactive was incorporated into a chitosan solution, which was then cross-linked with sodium tripolyphosphate (TPP) to form nanoparticles. Particle size and morphology were characterized using dynamic light scattering (DLS) and scanning electron microscopy (SEM).

Inclusion Complexes:

β-Cyclodextrin inclusion complexes were formed by mixing the herbal bioactives with β-cyclodextrin in an aqueous medium under constant stirring. The mixture was then lyophilized to obtain a stable complex, which was characterized using Fourier-transform infrared spectroscopy (FTIR) and X-ray diffraction (XRD) to confirm the formation of the inclusion complex.

III. DISSOLUTION STUDIES

A comprehensive set of in vitro dissolution studies was conducted to assess the performance of various formulations integrating natural polymers. Using a USP dissolution apparatus maintained at 37 °C to simulate physiological conditions, the release profiles of the herbal bioactives were monitored over time. Formulations prepared as solid dispersions, nanoparticles, and inclusion complexes were evaluated, with high-performance liquid chromatography (HPLC) employed to accurately quantify the concentration of the active compounds in the dissolution medium. The results revealed a significant enhancement in the dissolution rates of the polymer-based formulations compared to the pure herbal extracts. This improvement was largely attributed to the increased wettability, reduced particle size,

and amorphization of the bioactives, which collectively facilitated enhanced water penetration and solubilization. Additionally, the controlled release observed in these systems suggested that the choice of polymer not only affected the initial burst release but also sustained the release over an extended period. Overall, these studies underscore the critical role of natural polymers in optimizing the dissolution and bioavailability of poorly soluble herbal bioactives.

IV. RESULTS AND DISCUSSION

Table 1. Dissolution Profile of Herbal Bioactives in Various Formulations

Bioactive	Formulation Type	Polymer Used	% Dissolution at 30 min	% Dissolution at 60 min	Remarks
Curcumin	Pure	–	20%	30%	Poor dissolution
Curcumin	Solid Dispersion	HPMC	65%	85%	Improved wettability and amorphization
Curcumin	Nanoparticles	Chitosan	75%	90%	Highest enhancement; rapid release
Curcumin	Inclusion Complex	β-Cyclodextrin	70%	88%	Moderate improvement; reduced crystallinity
Quercetin	Pure	–	25%	35%	Poor dissolution
Quercetin	Solid Dispersion	HPMC	60%	80%	Significant improvement
Quercetin	Nanoparticles	Chitosan	68%	85%	Good enhancement; increased surface area
Quercetin	Inclusion Complex	β-Cyclodextrin	65%	82%	Moderate improvement
Resveratrol	Pure	–	22%	32%	Poor dissolution

Resveratrol	Solid Dispersion	HPMC	58%	78%	Improved solubility
Resveratrol	Nanoparticles	Chitosan	70%	88%	Enhanced dissolution; high efficiency
Resveratrol	Inclusion Complex	β-Cyclodextrin	66%	84%	Moderate improvement

These results underscore the critical role of natural polymers in overcoming the solubility challenges of herbal bioactives. The optimized formulations, particularly the nanoparticulate systems, not only demonstrated rapid initial release but also maintained a sustained release profile. This dual enhancement is vital for achieving consistent therapeutic levels and maximizing the bioavailability of these compounds. The observed improvements across different herbal bioactives confirm the broad applicability of these polymer-based strategies in the field of phytopharmaceuticals.

The formulation innovations demonstrated a significant enhancement in the dissolution rates of all three herbal bioactives. Solid dispersions with HPMC resulted in an amorphous state of the bioactives, which contributed to improved wettability and rapid dissolution. Chitosan-based nanoparticles provided a high surface area-to-volume ratio, thereby facilitating faster dissolution and enhanced bioavailability. In the case of β-cyclodextrin inclusion complexes, the encapsulation of the hydrophobic molecules led to a notable increase in aqueous solubility, as evidenced by the marked reduction in crystallinity observed in XRD patterns.

Each formulation approach exhibited distinct advantages. The solid dispersion technique was simple and scalable, making it suitable for industrial applications. Nanoparticulate systems, while offering enhanced bioavailability, required careful optimization of particle size and surface charge to maintain stability. Inclusion complexation, on the other hand, provided the dual benefits of solubility enhancement and protection against chemical degradation, albeit with the necessity for rigorous control over complexation conditions.

The improved dissolution profiles are attributed to the interactions between natural polymers and phytoconstituents. These interactions disrupt the crystalline lattice of the bioactives, facilitating a transition to an amorphous form that is more soluble in aqueous environments. Moreover, the hydrophilic nature of the polymers enhances water penetration and wettability, which further contributes to the dissolution process.

The study underscores the potential of natural polymers as versatile tools for formulation innovations in herbal medicine. By strategically leveraging polymer-phytoconstituent

interactions, it is possible to develop more effective and patient-friendly herbal formulations that overcome the longstanding challenge of poor solubility.

V. CONCLUSION

The research highlights that innovative formulation strategies employing natural polymers can significantly boost the dissolution of herbal bioactives. Through techniques such as solid dispersions, nanoparticle fabrication, and inclusion complexation, the inherent limitations of poorly soluble plant-derived compounds can be effectively mitigated. These advancements not only enhance the therapeutic efficacy of herbal medicines but also pave the way for more sustainable and biocompatible drug delivery systems. Future work should focus on in vivo validation of these formulations and the exploration of additional natural polymers to further optimize herbal bioavailability.

VI. REFERENCES

1. Sharma, P., Mehta, M., & Dhanawat, M. (2020). Chitosan-based nanocarriers for enhanced solubility of curcumin. Journal of Drug Delivery Science and Technology, 57, 101109.

2. Gupta, M., & Mishra, B. (2019). Cyclodextrin complexes in herbal formulations: Improving solubility and stability. Phytotherapy Research, 33(4), 785–793.

3. Patel, R., Shah, D., & Thakkar, V. (2021). Solid dispersions as a strategy to improve the solubility of phytoconstituents. International Journal of Pharmaceutics, 601, 120566.

4. Singh, R., & Lillard, J.W. (2009). Nanoparticle-based targeted drug delivery: Principles and clinical applications. Experimental and Molecular Pathology, 86(3), 215–223.

5. Das, S., Ng, K.Y., & Kanaujia, P. (2012). Enhancing the bioavailability of poorly soluble drugs via solid dispersion techniques. International Journal of Pharmaceutics, 428(1–2), 1–9.

6. Goyal, A.K., & Rath, G. (2013). Advances in herbal drug delivery systems: Innovations and future trends. Recent Patents on Drug Delivery & Formulation, 7(1), 1–8.

7. Rajinikanth, P.S., & Mishra, B. (2008). Intranasal delivery of herbal actives using alginate microspheres. Drug Delivery, 15(6), 331–339.

8. Khadka, P., et al. (2014). Pharmaceutical particle technologies: A review on improving drug solubility, dissolution, and bioavailability. Asian Journal of Pharmaceutical Sciences, 9(6), 304–316.

9. Kaur, L.P., & Guleri, T.K. (2013). Recent trends in topical gel formulations for improved bioavailability. Asian Journal of Biomedical and Pharmaceutical Sciences, 3(17), 1–5.

10. Pandey, A., Tripathi, S.K., & Rai, S.K. (2020). Biopolymer-based nanoformulations for enhancing the solubility of phytochemicals: A comprehensive review. Current Drug Delivery, 17(10), 881–894.